



Contents lists available at ScienceDirect

Molecular Phylogenetics and Evolution

journal homepage: www.elsevier.com/locate/ympevPhylogeny of *Nolana* (Solanaceae) of the Atacama and Peruvian deserts inferred from sequences of four plastid markers and the nuclear *LEAFY* second intronTieyao Tu^{a,b,c}, Michael O. Dillon^d, Hang Sun^a, Jun Wen^{a,e,*}^a Key Laboratory of Biodiversity and Biogeography, Kunming Institute of Botany, Chinese Academy of Sciences, Kunming, Yunnan 650204, PR China^b Graduate School of Chinese Academy of Sciences, Beijing 100039, PR China^c South China Botanical Garden, The Chinese Academy of Sciences, Guangzhou 510650, PR China^d Department of Botany, The Field Museum, 1400 South Lake Shore Drive, Chicago, IL 60605, USA^e Department of Botany, MRC-166, National Museum of Natural History, Smithsonian Institution, P.O. Box 37012, Washington, DC 20013-7012, USA

ARTICLE INFO

Article history:

Received 18 March 2008

Revised 1 July 2008

Accepted 26 July 2008

Available online xxxxx

Keywords:

Gene duplication

Lineage sorting

*ndhF**psbA-trnH*

Reticulate evolution

*rps16-trnK**trnC-psbM*

ABSTRACT

The phylogeny of *Nolana* (Solanaceae), a genus primarily distributed in the coastal Atacama and Peruvian deserts with a few species in the Andes and one species endemic to the Galápagos Islands, was reconstructed using sequences of four plastid regions (*ndhF*, *psbA-trnH*, *rps16-trnK* and *trnC-psbM*) and the nuclear *LEAFY* second intron. The monophyly of *Nolana* was strongly supported by all molecular data. The *LEAFY* data suggested that the Chilean species, including *Nolana sessiliflora*, the *N. acuminata* group and at least some members of the *Alona* group, are basally diverged, supporting the Chilean origin of the genus. Three well-supported clades in the *LEAFY* tree were corroborated by the SINE (short interspersed elements) or SINE-like insertions. Taxa from Peru are grouped roughly into two clades. *Nolana galapagensis* from the Galápagos Island is most likely to have derived from a Peruvian ancestor. The monophyly of the morphologically well-diagnosed *Nolana acuminata* group (*N. acuminata*, *N. baccata*, *N. paradoxa*, *N. parviflora*, *N. pterocarpa*, *N. rupicola* and *N. elegans*) was supported by both plastid and *LEAFY* data. Incongruence between the plastid and the *LEAFY* data was detected concerning primarily the positions of *N. sessiliflora*, *N. galapagensis*, taxa of the *Alona* group and the two Peruvian clades. Such incongruence may be due to reticulate evolution or in some cases lineage sorting of plastid DNA. Incongruence between our previous GBSSI trees and the plastid-*LEAFY* trees was also detected concerning two well-supported major clades in the GBSSI tree. Duplication of the GBSSI gene may have contributed to this incongruence.

© 2008 Published by Elsevier Inc.

1. Introduction

The genus *Nolana* L. f. consists of 89 species primarily distributed in the Atacama and Peruvian deserts, with 43 species in Peru, 49 species in Chile, a few species in the inland regions of the Andes (e.g., *N. chapiensis*, *N. lezamae*, *N. sessiliflora*, *N. urubambae*, *N. tarapacana*) and one species endemic to the Galápagos Islands. It is the fourth largest genus in the family Solanaceae after *Solanum* (ca. 1500 species), *Cestrum* (ca. 160 species) and *Physalis* (ca. 120 species). Members of this genus are annuals, or perennial herbs or woody shrubs. Adapting to the unusual arid lomas environment in coastal Peru and Chile (see Dillon et al., 2007), species of *Nolana* have developed somewhat succulent leaves arranged in rosettes and shoots with short internodes. When the water conditions are

favorable for growing, the rosettes may increase in size and the flowering period may prolong. This suite of adaptive characters may confound phylogenetic analysis using morphology. Flowers of *Nolana* are hermaphroditic with corollas varying greatly in size, actinomorphic to weakly zygomorphic, tubular-salverform to campanulate and white to blue in color with variable colored spots and veins in the throat. The most significant character separating *Nolana* from other Solanaceae taxa is the presence of the unusual sclerified fruits called mericarps in this genus (Knapp, 2002). The mericarp number can be reduced to as few as two or as many as 30, often with several seeded mericarps arising through incomplete radial fission of the fertile carpels (Bondeson, 1896; Saunders, 1936; Tago-Nakawaza and Dillon, 1999).

Because of its unique fruit type, *Nolana* has been widely accepted as a highly distinct group since its description by Linnaeus f. in 1762 (Don, 1838; Hunziker, 2001; Johnston, 1936; Mesa, 1981). The monophyly of *Nolana* is also strongly supported by sequence data from the plastid *matK* gene, the nuclear ribosomal internal transcribed spacer (ITS) (Tago-Nakawaza and Dillon, 1999) and partial

* Corresponding author. Address: Department of Botany, MRC-166, National Museum of Natural History, Smithsonian Institution, P.O. Box 37012, Washington, DC 20013-7012, USA. Fax: +1 202 786 2563.

E-mail address: Wenj@si.edu (J. Wen).

sequences of the nuclear granule-bound starch synthase I (GBSSI) gene (Dillon et al., 2007). Some workers recognize *Nolana* at the familial rank (Cronquist, 1981; Hunziker, 2001), or as a subfamily, i.e., Nolanoideae of Solanaceae (D'Arcy, 1979, 1991; Dahlgren, 1980; Takhtajan, 1997; Thorne, 1983). Data from plastid DNA restriction site mapping and plastid *ndhF* gene sequences, with most Solanaceous genera sampled, have strongly supported the placement of *Nolana* within the Solanaceae, and suggested its sister relationship with the tribe Lycieae (Olmstead and Palmer, 1992; Olmstead and Sweere, 1994; Olmstead et al., 1999).

In our previous efforts, the plastid *matK*, ITS (Tago-Nakawaza and Dillon, 1999) and the nuclear GBSSI sequences (Dillon et al., 2007) were employed to elucidate interspecific relationships of *Nolana*. The initial phylogenetic study using ITS and *matK* data sampling 37 species produced poorly resolved phylogenies. The sequence data from the third to the eighth exon of the GBSSI gene produced a much better resolved phylogeny of the genus. The GBSSI tree suggested the sister relationship between *N. sessiliflora* and the remainder of the genus, two strongly supported major clades for the remaining species and eight strongly to moderately supported subclades within the two major clades. The subclade (the *Nolana acuminata* group) comprised of *Nolana paradoxa*, *N. acuminata*, *N. reichei*, *N. elegans*, *N. rupicola*, *N. pterocarpa*, *N. baccata* and *N. parviflora* was supported by the GBSSI data. This subclade in the GBSSI tree is also supported by morphology and distribution, as all taxa in the subclade share the characters of basal rosettes, large showy flowers and 10–20 mericarps and are generally distributed in coastal Chile. The other seven subclades did not contradict relationships inferred from morphology and geographic distribution. However, the interspecific relationships within most subclades were largely unresolved. Furthermore, each of the two major clades includes species with diverse morphological characters and each has species from both Chile and Peru. Several mechanisms including adaptive radiation, reticulate evolution or gene duplication may lead to a clade of taxa with diverse morphology based on molecules. If the last scenario is true, the phylogeny based on these sequences may be misleading. Special caution should be made when using low-copy nuclear genes because they are prone to gene duplications through polyploidy or retrotransposition even losing gene copies (Sang, 2002). Duplication of the GBSSI gene has been reported in some groups of flowering plants, including Rosaceae (Evans et al., 2000), *Viburnum* (Carprifoliaceae) (Winkworth and Donoghue, 2004) and *Spartina* (Poaceae) (Fortune et al., 2007). In Solanaceae, initial phylogenetic studies detected only one copy of the GBSSI gene, such as in the diploid *Solanum* (Levin et al., 2006, 2005; Peralta and Spooner, 2001), the Iochrominae group of the tribe Physaleae (Smith and Baum, 2006), *Schizanthus* (Perez et al., 2006) and *Nolana*'s close relative Lycieae (Levin and Miller, 2005). In a recent study, more than two copies of this gene were found in Hyoscyameae, a polyploid group from the northern hemisphere closely related to *Nolana* (Yuan et al., 2006). Evolution of the GBSSI gene in Solanaceae thus may be more complex than previously thought, and the GBSSI phylogeny of Solanaceae taxa needs to be tested using additional markers.

In this study, we employed four plastid markers to test the phylogeny of *Nolana*. We chose *ndhF* because the gene has been used for a broad range of taxa in Solanaceae (Olmstead and Sweere, 1994). Partial sequences between the *trnC* and the *psbM* genes were sequenced because of its relatively high rate of nucleotide substitution in *Panax* (Araliaceae) (Lee and Wen, 2004). The intergenic region *trnH-psbA* (Shaw et al., 2005) has been demonstrated to be highly variable at intraspecific and interspecific levels in the Solanaceous genus *Petunia* (Lorenz-Lemke et al., 2006). Considerable variation of the *rps16-trnK* spacer (Shaw et al., 2007) was detected from the alignment among sequences of *Solanum*, *Nicotiana* and *Atropa* (GenBank Accession Nos. NC007500, NC001879, NC008096, NC007943, NC004561).

We also used the nuclear *LEAFY* gene in our analysis. The *LEAFY* gene is a homeotic gene which regulates the floral meristem induction during the early stages of reproductive ontogeny (Blazquez, 1997; Blazquez et al., 1997; Schultz and Haughn, 1991; Wada et al., 2002; Weigel, 1995). In some cases, it affects the vegetative morphogenesis (Hofer et al., 1997; Kelly et al., 1995; Pouteau et al., 1997). It was first described as *FLORICAULA* in *Antirrhinum majus* (Coen et al., 1990) and then as *LEAFY* in *Arabidopsis thaliana* (Schultz and Haughn, 1991). Some other names have been used to designate the orthologs of *LEAFY* in plants, such as *NFL* in *Nicotiana tabacum* (Kelly et al., 1995), *Imp-flo* in *Impatiens* (Pouteau et al., 1997) and *alf* in *Petunia* (Souer et al., 1998). More than one copy has been reported for the *LEAFY* orthologs in the gymnosperms, and some basal or polyploid angiosperms (Bombliet et al., 2003; Cronk, 2001; Frohlich and Meyerowitz, 1997; Frohlich and Parker, 2000; Theissen, 2000). Whereas this gene has been generally suggested to be single-copy in most diploid angiosperm species studied so far, exceptions include two or more possible copies in the diploid *Eucalyptus* L. (Southerton et al., 1998) and at least two clear copies in certain taxa of the Lamiales (Aagaard et al., 2005, 2006), Leguminosae (Archambault and Bruneau, 2004) and Brassicaceae (Baum et al., 2005). In Solanaceae, only one copy of *alf*, the ortholog of *LEAFY*, was presumed for *Petunia* Juss. based on the southern blot and the inflorescence cDNA library screening experiment (Souer et al., 1998). Two copies of *NFL*, the homolog of *FLORICAULA* and *LEAFY*, were detected in the cultivated allotetraploid *Nicotiana tabacum* L. As expected, a single-copy of this gene was observed from both paternal (*N. sylvestris* Speg.) and maternal (*N. tomentosiformis* Goodspeed) parents of the allotetraploid *N. tabacum* (Kelly et al., 1995). The generally low-copy number of *LEAFY* in angiosperms and the relatively high level of variation within the introns make it an excellent candidate as a phylogenetic marker for resolving interspecific even intraspecific relationships or for testing hypothesis of hybridization (Grob et al., 2004; Hoot and Taylor, 2001; Howarth and Baum, 2005; Oh and Potter, 2003, 2005). The first use of *LEAFY* for phylogenetic study of Solanaceae (Smith and Baum, 2006) demonstrated that the second intron of *LEAFY* contains more informative characters than those from ITS and GBSSI together. The *LEAFY* sequences were also shown to be useful in detecting hybridization in the Iochrominae group of Solanaceae.

Objectives of this study are to: (1) elucidate the interspecific relationships within *Nolana* using multiple molecular markers; (2) test the GBSSI phylogeny of the genus and (3) evaluate the phylogenetic utility of the nuclear *LEAFY* second intron.

2. Materials and methods

2.1. Taxon sampling, DNA extraction and amplification and sequencing

All 63 species analyzed in our previous GBSSI study (Dillon et al., 2007) as well as two additional species, *Nolana tocopilensis* and *N. ivaniana*, were sequenced for four plastid markers in the present study. However, only 55 species were sequenced for the nuclear *LEAFY* gene because of difficulties in amplifying this gene from some degenerated leaf tissue samples. DNA extractions followed Dillon et al. (2007) and voucher information was presented in Table 1.

Target regions were amplified in 25 μ l reaction-mixture volumes using the Bioline Taq polymerase and associated reagents at 2.0 mM MgCl₂ concentration except for *trnC-psbM*, which used 4.0 mM MgCl₂. Primers for *ndhF*, *trnH-psbA* and *rps16-trnK* followed Olmstead and Sweere (1994), Shaw et al. (2005, 2007), respectively. The primers *trnC* (5'-CCAGTTCAAATCCGGGTGTC-3') and 2039R (5'-TTTCTACTTATCATTTACG-3') were used to amplify the *trnC-psbM*

Table 1
List of the taxa sampled with geographic origins, voucher numbers and GenBank accession numbers

| Species | Location | Voucher | GenBank Accession No. | | | | |
|--|-----------------------------|---------------------------|-----------------------|------------------|-------------------|------------------|--|
| | | | <i>ndhF</i> | <i>psbA-trnH</i> | <i>rps16-trnK</i> | <i>trnC-psbM</i> | <i>LEAFY</i> |
| <i>Grabowskia glauca</i> (Phil.) I.M. Johnst. | Chile (Antofagasta) | Dillon 8581 (F) | EU742303 | EU742439 | EU742371 | EU742507 | — |
| <i>Lycium americana</i> Jacq. | Peru (Arequipa) | Quipuscoa 2862 (F) | EU742304 | EU742440 | EU742372 | EU742508 | EU742190 |
| <i>L. deserti</i> Phil. | Chile (Antofagasta) | Dillon 8545 (F) | EU742305 | EU742441 | EU742373 | EU742509 | — |
| <i>Nolana acuminata</i> (Miers) Miers ex Dunal | Chile (Región II) | Dillon 8100 (F) | EU742307 | EU742443 | EU742375 | EU742511 | — |
| <i>N. adansonii</i> (Roem. & Schult.) I.M. Johnst. | Peru (Arequipa) | Dillon 8984 (F) | EU742308 | EU742444 | EU742376 | EU742512 | EU742223 clone3 EU742224 clone7 EU742225 clone4 EU742226 clone8 |
| <i>N. albescens</i> (Phil.) I.M. Johnst. | Chile (Región III) | Dillon 8666 (F) | EU742309 | EU742445 | EU742377 | EU742513 | — |
| <i>N. aplocaryoides</i> (Gaudich.) I.M. Johnst. | Chile (Región II) | Thompson 01 (F) | EU742310 | EU742446 | EU742378 | EU742514 | EU742192 |
| <i>N. arenicola</i> I.M. Johnst. | Peru (Arequipa) | Dillon 8763 (F) | EU742311 | EU742447 | EU742379 | EU742515 | EU742275 clone3 EU742276 clone1 EU742279 clone15 |
| <i>N. arequipensis</i> M.O. Dillon & Quipuscoa | Peru (Arequipa) | Dillon 8790 (F) | EU742312 | EU742448 | EU742380 | EU742516 | EU742241 EU742242 |
| <i>N. aticoana</i> Ferreyra | Peru (Arequipa) | Dillon 8806 (F) | EU742313 | EU742449 | EU742381 | EU742517 | EU742254 |
| <i>N. baccata</i> (Lindl.) Dunal | Chile (Región III) | Dillon 8612 (F) | EU742314 | EU742450 | EU742382 | EU742518 | EU742297 |
| <i>N. balsamiflua</i> (Gaudich.) Mesa | Chile (Región II) | Dillon 5726 (F) | EU742315 | EU742451 | EU742383 | EU742519 | EU742280 clone6 EU742281 clone2 EU742282 clone4 |
| <i>N. carnosa</i> (Lindl.) Miers ex Dunal | Chile (Caldera) | Teillier & Dillon s.n. | EU742316 | EU742452 | EU742384 | EU742520 | EU742287 |
| <i>N. cerrateana</i> Ferreyra | Peru (Arequipa) | Quipuscoa 2890 (F) | EU742317 | EU742453 | EU742385 | EU742521 | EU742236 clone4 EU742237 clone5 EU742238 clone7 EU742252 clone1 |
| <i>N. chancoana</i> M.O. Dillon & Quipuscoa | Peru (Arequipa) | Dillon 8789 (F) | EU742318 | EU742454 | EU742386 | EU742522 | EU742263 clone1 EU742264 clone5 EU742265 clone4 |
| <i>N. chapiensis</i> M.O. Dillon & Quipuscoa | Peru (Arequipa) | Quipuscoa 2785 (F) | EU742319 | EU742455 | EU742387 | EU742523 | EU742261 |
| <i>N. clivicola</i> (I.M. Johnst.) I.M. Johnst. | Chile (Región II) | Dillon 8584 (F) | EU742320 | EU742456 | EU742388 | EU742524 | EU742259 |
| <i>N. coelestis</i> (Lindl.) Miers ex Dunal | Chile (Región IV) | Miller 0498 (US) | EU742321 | EU742457 | EU742389 | EU742525 | EU742289 |
| <i>N. confinis</i> I.M. Johnst. | Peru (Moquegua) | Quipuscoa 2921A (F) | EU742322 | EU742458 | EU742390 | EU742526 | EU742249 clone3 EU742250 clone1 EU742251 clone2 |
| <i>N. crassulifolia</i> Poepp. | Chile (Región III) | Dillon 8744 (F) | EU742323 | EU742459 | EU742391 | EU742527 | EU742204 |
| <i>N. diffusa</i> I.M. Johnst. | Chile (Región II) | Dillon 8076 (F) | EU742324 | EU742460 | EU742392 | EU742528 | EU742195 clone6 EU742197 clone4 EU742198 clone5 EU742199 clone3 |
| <i>N. divaricata</i> (Lindl.) I.M. Johnst. | Chile (Región II) | Dillon 5666 (F) | EU742325 | EU742461 | EU742393 | EU742529 | EU742205 |
| <i>N. elegans</i> (Phil.) Reiche | Chile (Región II) | Dillon 8071 (F) | EU742326 | EU742462 | EU742394 | EU742530 | — |
| <i>N. filifolia</i> (Hook. & Arn.) I.M. Johnst. | Chile (Región III) | Dillon 8655 (F) | EU742327 | EU742463 | EU742395 | EU742531 | EU742288 |
| <i>N. flaccida</i> (Phil.) I.M. Johnst. | Chile (Región II) | Dillon 8101 (F) | EU742328 | EU742464 | EU742396 | EU742532 | — |
| <i>N. galapagensis</i> (Christoph.) I.M. Johnst. | Ecuador (Galápagos Islands) | Dillon 8504 (F) | EU742329 | EU742465 | EU742387 | EU742533 | EU742268 |
| <i>N. gayana</i> (Gaudich.) Koch | Peru (Ancash) | Leiva 2603 (F) | EU742330 | EU742466 | EU742398 | EU742534 | EU742256 |
| <i>N. glauca</i> (I.M. Johnst.) I.M. Johnst. | Chile (Chañaral) | Luebert & Becker 2841 (F) | EU742331 | EU742467 | EU742399 | EU742535 | — |
| <i>N. humifusa</i> (Gouan) I.M. Johnst. | Peru | Freyre s.n. (F) | EU742332 | EU742468 | EU742400 | EU742536 | EU742257 clone5 EU742258 clone12 |
| <i>N. incana</i> (Phil.) I.M. Johnst. | Chile (Región II) | Dillon 8072 (F) | EU742333 | EU742469 | EU742401 | EU742537 | EU742206 clone2 EU742207 clone8 EU742208 clone5 |
| <i>N. inflata</i> Ruiz & Pav. | Peru (Arequipa) | Dillon 8805 (F) | EU742334 | EU742470 | EU742402 | EU742538 | EU742266 |
| <i>N. intonsa</i> I.M. Johnst. | Chile (Región I) | Dillon & Finger 8590 (F) | EU742335 | EU742471 | EU742403 | EU742539 | EU742227 clone3 |

(continued on next page)

Table 1 (continued)

| Species | Location | Voucher | GenBank Accession No. | | | | |
|---|---------------------|----------------------------|-----------------------|------------------|-------------------|------------------|------------------|
| | | | <i>ndhF</i> | <i>psbA-trnH</i> | <i>rps16-trnK</i> | <i>trnC-psbM</i> | <i>LEAFY</i> |
| <i>N. ivaniana</i> Ferreyra | Peru (Arequipa) | Dillon 8973 (F) | EU742336 | EU742472 | EU742404 | EU742540 | EU742228 clone4 |
| | | | | | | | EU742229 clone1 |
| | | | | | | | EU742230 clone2 |
| | | | | | | | EU742231 clone5 |
| | | | | | | | EU742269 clone2 |
| <i>N. johnstonii</i> Vargas | Peru (Moquega) | Quipuscoa 2919 (F) | EU742337 | EU742473 | EU742405 | EU742541 | EU742270 clone1 |
| | | | | | | | EU742271 clone4 |
| | | | | | | | EU742272 clone5 |
| | | | | | | | EU742253 |
| | | | | | | | EU742216 |
| | | | | | | | EU742262 |
| | | | | | | | EU742211 |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. lachimbensis</i> M.O. Dillon & Luebert | Chile (Región II) | Dillon 8591 (F) | EU742338 | EU742474 | EU742406 | EU742542 | EU742232 clone3 |
| | | | | | | | EU742233 clone2 |
| | | | | | | | EU742234 clone1 |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. laxa</i> (Miers) I.M. Johnst. | Peru (Lima) | Freyre s.n. | EU742339 | EU742475 | EU742407 | EU742543 | EU742239 clone4 |
| | | | | | | | EU742240 clone7 |
| | | | | | | | EU742248 clone5 |
| | | | | | | | EU742290 |
| | | | | | | | EU742293 |
| | | | | | | | EU742209 |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. leptophylla</i> (Miers) I.M. Johnst. | Chile (Región II) | Dillon 8181 (F) | EU742340 | EU742476 | EU742408 | EU742544 | EU742295 clone4 |
| | | | | | | | EU742296 clone2 |
| | | | | | | | EU742298 clone1 |
| | | | | | | | EU742196 |
| | | | | | | | EU742285 clone1 |
| | | | | | | | EU742286 clone2 |
| | | | | | | | EU742291 clone2 |
| | | | | | | | EU742292 clone1 |
| | | | | | | | EU742194 |
| | | | | | | | EU742273 clone4 |
| <i>N. lezamae</i> M.O. Dillon, S. Leiva & Quipuscoa | Peru (Ancash) | Leiva 2212 (F) | EU742341 | EU742477 | EU742409 | EU742545 | EU742274 clone1 |
| | | | | | | | EU742277 clone3 |
| | | | | | | | EU742278 clone7 |
| | | | | | | | EU742210 clone3 |
| | | | | | | | EU742212 clone1 |
| | | | | | | | EU742213 clone5 |
| | | | | | | | EU742214 clone4 |
| | | | | | | | EU742215 clone2 |
| | | | | | | | EU742283 clone10 |
| | | | | | | | EU742284 clone19 |
| <i>N. linearifolia</i> Phil. | Chile (Región II) | Dillon 8585A (F) | EU742342 | EU742478 | EU742410 | EU742546 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. lycioides</i> I.M. Johnst. | Peru (Arequipa) | Quipuscoa 2913 (F) | EU742343 | EU742479 | EU742411 | EU742547 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. mollis</i> Phil. | Chile (Antofagasta) | Luebert & Gracia 2757 (F) | EU742344 | EU742480 | EU742412 | EU742548 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. pallida</i> I.M. Johnst., | Peru (Arequipa) | Dillon 8938 (F) | EU742345 | EU742481 | EU742413 | EU742549 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. paradoxa</i> Lindl. | Chile (IV) | Dillon 8686 (F) | EU742346 | EU742482 | EU742414 | EU742550 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. parviflora</i> (Phil.) Phil. | Chile (Región II) | Luebert 2895 (F) | EU742347 | EU742483 | EU742415 | EU742551 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. peruviana</i> (Gaudich.) I.M. Johnst. | Chile (Región II) | Luebert 2559 (F) | EU742348 | EU742484 | EU742416 | EU742552 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. peruviana</i> (Gaudich.) I.M. Johnst. | Chile (Región II) | Dillon 8567 (F) | EU742349 | EU742485 | EU742417 | EU742553 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. plicata</i> I.M. Johnst. | Peru (Arequipa) | Dillon 8782 (F) | EU742350 | EU742486 | EU742418 | EU742554 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. pterocarpa</i> Phil. | Chile (Región III) | Teillier & Dillon 4926 (F) | EU742351 | EU742487 | EU742419 | EU742555 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. ramosissima</i> I.M. Johnst. | Chile (Región II) | Dillon 8085 (F) | EU742352 | EU742488 | EU742920 | EU742556 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. rostrata</i> (Lindl.) Miers ex Dunal | Chile (Región III) | Dillon 8615B (F) | EU742353 | EU742489 | EU742921 | EU742557 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. rupicola</i> Gaudich. | Chile (Región II) | Dillon 8647 (F) | EU742354 | EU742490 | EU742922 | EU742558 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. salsoloides</i> (Lindl.) I.M. Johnst. | Chile (Región II) | Dillon 8584A (F) | EU742355 | EU742491 | EU742923 | EU742559 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. scaposa</i> Ferreyre | Peru (Arequipa) | Dillon 8940 (F) | EU742356 | EU742492 | EU742924 | EU742560 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. sedifolia</i> Poepp. | Chile (Región II) | Dillon 8721 (F) | EU742357 | EU742493 | EU742925 | EU742561 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. sessiliflora</i> Phil. | Chile (Región II) | Dillon 8644A (F) | EU742358 | EU742494 | EU742926 | EU742562 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. spathulata</i> Ruiz & Pav. | Peru (Arequipa) | Quipuscoa 2885 (F) | EU742359 | EU742495 | EU742927 | EU742563 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. sphaerophylla</i> (Phil.) Mesa ex Dillon | Chile (Chañaral) | Luebert & Becker 2840 (F) | EU742360 | EU742496 | EU742928 | EU742564 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| <i>N. stenophylla</i> I.M. Johnst. | Chile (Región II) | Dillon 5909 (F) | EU742361 | EU742497 | EU742929 | EU742565 | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |
| | | | | | | | – |

| | | | | | | | |
|--|-------------------|--------------------|----------|----------|----------|----------|-----------------|
| <i>N. tarapacana</i> (Phil.) I.M. Johnst. | Chile (Región I) | Teillier 4803 (F) | EU742362 | EU742498 | EU742930 | EU742566 | — |
| <i>N. thinothila</i> I.M. Johnst. | Peru (Arequipa) | Quipuscoa 2918 (F) | EU742363 | EU742499 | EU742931 | EU742567 | EU742222 |
| <i>N. tocopillensis</i> (Phil.) I.M. Johnst. | 7 | Dillon 8586 (F) | EU742364 | EU742500 | EU742932 | EU742568 | EU742193 |
| <i>N. tomentella</i> Ferreyra | Peru (Arequipa) | Dillon 8784 (F) | EU742365 | EU742501 | EU742433 | EU742569 | EU742243 clone7 |
| <i>N. urubamba</i> Vargas | Peru (Cusco) | Tupayachi 3858 (F) | EU742366 | EU742502 | EU742434 | EU742570 | EU742244 clone6 |
| <i>N. villosa</i> (Phil.) I.M. Johnst. | Chile (Región II) | Dillon 8074 (F) | EU742367 | EU742503 | EU742435 | EU742571 | EU742245 clone3 |
| <i>N. volcanica</i> Ferreyra | Peru (Arequipa) | Quipuscoa 2930 (F) | EU742368 | EU742504 | EU742436 | EU742572 | EU742246 clone4 |
| <i>N. weissiana</i> Ferreyra | Peru (Arequipa) | Dillon 8898 (F) | EU742369 | EU742505 | EU742437 | EU742573 | EU742247 clone1 |
| <i>N. wenderrmannii</i> I.M. Johnst. | Chile (Región IV) | Dillon 8665 (F) | EU742370 | EU742506 | EU742438 | EU742574 | EU742217 clone1 |
| <i>Phrodus microphylla</i> (Miers) | Chile (Región II) | Dillon 8643 (F) | EU742306 | EU742442 | EU742374 | EU742510 | EU742218 clone4 |
| | | | | | | | EU742219 clone5 |
| | | | | | | | EU742220 clone2 |
| | | | | | | | EU742221 clone3 |
| | | | | | | | EU742191 |

region and one internal primer 690F (5'-TTTATATTATAGAGA TAGGGGAC-3') was designed for sequencing.

The second intron of the *LEAFY* gene was initially amplified and sequenced from a subset of taxa using degenerate primers F2 and R1 (Howarth and Baum, 2005). These sequences were used to design *Nolana* specific primers (LFYNo13F: 5'-TATTGCCAAGGAACGA GGTG-3' and LFYNo13R: 5'-CGTACCTGAACACTTGATTG-3'). Two internal sequencing primers were also designed (LFYNo15F: 5'-TACGGACTGATGGGCTGAAC-3' and LFYNo15R: 5'-GACAAGTTACA GGTGGAGATAC-3'). Most amplified products contained one band and were sequenced directly. Cloning was conducted using the TOPO TA cloning kit (Invitrogen, Carlsbad, CA, USA) when ambiguous sequences were obtained by direct sequencing or when more than one band was detected during the amplification. At least five clones representing each band of the PCR products were sequenced. To capture potential hidden copies, we selected four samples which showed multiple bands, and used low annealing temperature (45 °C) in the PCR reactions and sequenced 20 clones for each of the four samples.

The PCR reactions for *LEAFY* differed from the *ndhF* reactions in that 10 μM BSA was used in the *LEAFY* amplification. The PCR program for the *LEAFY* amplification was 95 °C for 3 min, then 35 cycles of 94 °C for 40 s, 50 °C for 40 s, 72 °C for 2 min, followed by a final extension of 72 °C for 10 min. The amplified products were then purified using the polyethylene glycol (PEG) precipitation.

Cycle sequencing was conducted using the BigDye 3.1 reagents with an ABI 3700 automated sequencer (Applied Biosystems, Foster City, CA, USA). The program Sequencher 4.5 (Gene Codes Corporation, 2005) was used to evaluate chromatograms for base confirmation and to edit contiguous sequences. Sequences were initially aligned with ClustalX version 1.83 (Thompson et al., 1997), followed by manual adjustments on Se-Al v2.0a11 (Rambaut, 2007).

2.2. Phylogenetic analyses

Parsimony analysis was performed using a heuristic search with 100 random sequences addition replicates, tree bisection-reconnection (TBR) swapping, collapse of zero-length branches, multiple tree option in effect and character state changes equally weighted in the analysis. Because too many trees were found for the *LEAFY* data, trees were limited to 10,000 during each of 10 random sequences addition replicate. Gaps were treated either as missing data or coded as simple indels using the program GapCoder (Young and Healy, 2003). Bootstrap values (BP) (Felsenstein, 1985) of the internal nodes were obtained with 500 replicates. In each replicate, we performed 10 random sequences addition replicates following by tree bisection-reconnection (TBR) swapping algorithm and keeping no more than 1000 trees per replicate.

Bayesian inference (Rannala and Yang, 1996) was conducted using MrBayes version 3.1.2 (Ronquist and Huelsenbeck, 2003) with the model estimated by Modeltest version 3.7 (Posada and Buckley, 2004; Posada and Crandall, 1998). The Markov chain Monte Carlo algorithm was run for 2,000,000 generations with four incrementally heated chains, starting from random trees and sampling one out of every 100 generations. The first 2000–5000 trees were discarded, depending on when chains appeared to have become stationary, and the remaining trees were used to construct the Bayesian consensus tree. Internodes with posterior probabilities (PP) ≥ 95% were considered statistically significant.

3. Results

3.1. Phylogenetic analyses of plastid DNA data

The four plastid markers had 5172 aligned positions, of which 220 were variable (4.2%) and 150 (2.9%) were parsimony-informa-

tive (PI). The aligned length of each marker was 1998 from the *ndhF* gene, 815 from *rps16-trnK*, 501 from *psbA-trnH* and 1858 from partial sequences of *trnC-psbM*. Treating gaps as missing data, the parsimony analysis generated 7717 equally most parsimonious trees (MPTs) with a tree length of 268 steps, a consistency index (CI) of 0.84, a consistency index excluding uninformative characters of 0.78 and a retention index (RI) of 0.95 (Table 2). The strict consensus tree is presented in 1. In the Bayesian analysis, 5000 trees were eliminated before generating the 50% majority-rule tree. The topology of the tree is similar to that of the MPTs. All nodes with high bootstrap value (>90%) had high PP (1.0) values as well. Some nodes with moderate to low bootstrap values also had good PP support, such as the cp-I clade, the cp-G clade, the cp-C clade and the cp-FH clade (Fig. 1).

Except for *ndhF*, all other three plastid DNA regions contained gaps. There were 25 new indel characters in the plastid DNA data set, of which ten were parsimony-informative, including eight repeats, one deletion and one insertion. All the unambiguous indels supported the topology of the base-substitution tree with two exceptions. One was the 7-base repeat in the outgroup and the cp-D clade, and the other was the 42-base repeat detected in the outgroup taxa *Grabowskia glauca* and *Phrodus microphylla*. The analysis of the plastid DNA with indels as new characters produced 15448 MPTs with a tree length of 296 steps, a CI of 0.85, a CI excluding uninformative characters of 0.78, and an RI of 0.95. The topology of the strict consensus tree from gaps as new characters was identical to that of the tree with gaps as missing data. The bootstrap values of clades were similar in both analyses.

3.2. Phylogenetic analysis of *LEAFY*

Amplification of the *LEAFY* second intron yielded one or two bands. All sequences from the larger bands can be aligned with *LEAFY* sequences of Solanaceae from GenBank. The sequences of the smaller bands did not match any *LEAFY* sequences or other genes. The nature of the smaller fragments remained unknown and these sequences were not included in the phylogenetic analysis.

Variation in the second intron of the *LEAFY* gene is higher than that in the four plastid markers (Table 2). Sequences across 113 accessions ranged from 843 to 1771 bp and had an aligned length of 4175 bp. Of these 4175 characters, 900 were variable (21%) and 564 were parsimony-informative (13.5%) (Table 2). Treating gaps as missing data, the parsimony analysis yielded 10,920 MPTs with a tree length of 1493 steps, a CI of 0.75, a CI excluding uninformative characters of 0.65, and an RI of 0.90. The strict consensus tree is presented in Fig. 2.

The indels in the second intron of the *LEAFY* gene composed of simple deletions, insertions, mononucleotide repeats or tandemly arranged multibase repeats. After the ambiguous blocks in the alignment were deleted, there were 325 indel characters, which ranged from 1 to 789 bp in size. The analysis treating indels as new characters had a tree length of 1898 steps, a CI of 0.74, a CI excluding uninformative characters of 0.65, and an RI of 0.90. The topology with indels as new characters was generally congruent with that of the tree when indels were treated as missing data. Nevertheless, the indel characters increased the bootstrap values of many clades (Fig. 2).

3.3. Phylogenetic results

The monophyly of *Nolana* has been recovered by both the plastid regions and sequences of the *LEAFY* second intron. Two large clades for *Nolana* (cp-I and cp-II) were detected in the plastid DNA tree, one containing taxa from Chile and the other with taxa from Chile and Peru. *Nolana acuminata*, *N. baccata*, *N. paradoxa*, *N.*

Table 2
Comparison of four plastid DNA markers and the nuclear *LEAFY* second intron

| Region | <i>ndhF</i> | <i>rps16-trnK</i> | <i>psbA-trnH</i> | <i>trnC-psbM</i> | Combined plastid DNA without gaps | Combined plastid DNA with gaps | <i>LEAFY</i> | <i>LEAFY</i> with gaps |
|---------------------------|-------------|-------------------|------------------|------------------|-----------------------------------|--------------------------------|--------------|------------------------|
| Aligned length | 1998 | 815 | 501 | 1858 | 5172 | 5196 | 4175 | 4500 |
| Variable sites/proportion | 76/3.8% | 40/4.9% | 46/9.1% | 58/3.1% | 220/4.2% | 245/4.7% | 900/21.0% | 1204/26.7% |
| PI sites/proportion | 52/2.6% | 25/3.1% | 28/5.6% | 45/2.4% | 150/2.9% | 162/3.1% | 564/13.5% | 752/16.7% |
| CI/RI | 0.94/0.98 | 0.95/0.99 | 0.75/0.91 | 0.89/0.97 | 0.84/0.95 | 0.85/0.95 | 0.75/0.90 | 0.74/0.90 |
| Tree length | 84 | 43 | 65 | 65 | 268 | 296 | 1493 | 1898 |

Note that the four plastid markers were sequenced for 68 OTUs and the *LEAFY* second intron had 113 OTUs. Note: PI, parsimony-informative; CI, consistency index; RI, retention index.




Fig. 1. Strict consensus tree of the most parsimonious trees of *Nolana* based on combined sequences of four chloroplast markers. Bootstrap values are provided above the branches leading to the nodes and Bayesian posterior probabilities are below the branches, bootstrap values below 50% and Bayesian values below 95% are not shown.

parviflora, *N. pterocarpa*, *N. rupicola*, *N. elegans*, *N. balsamiflua*, *N. linearifolia* and *N. sessiliflora* formed the cp-I clade, which was sister to the well-supported cp-II clade composed of the remaining species of the genus (Fig. 1). Within the cp-I clade, subclade cp-A consisting of *Nolana acuminata*, *N. baccata*, *N. paradoxa*, *N.*

parviflora, *N. pterocarpa*, *N. rupicola* and *N. elegans* was strongly supported, whereas the other clade (cp-D, generally corresponding to clade D in the previous GBSSI tree (Dillon et al., 2007)) including species of *N. balsamiflua*, *N. linearifolia* and *N. sessiliflora* is only weakly supported. In the cp-II clade, five subclades (cp-B,




Fig. 2. Strict consensus tree of the most parsimonious tree of *Nolana* based on sequences of the *LEAFY* second intron. Numbers next to the nodes indicate the bootstrap value based on the base-substitution data/the bootstrap values based on the analysis treating indels as new characters; The "-" means the bootstrap values are not changed. Bold branches indicate the nodes have Bayesian values > 95%. Clades are annotated as LFY-A to LFY-H. SINE1-SINE5 indicates the SINE or SINE-like insertions detected for the clades or species.

cp-C, cp-E, cp-G, cp-FH, corresponding to clade B, C, E, G and F and H in the previous GBSSI tree (Dillon et al., 2007) were recovered.

In the *LEAFY* tree, *N. sessiliflora* is sister to a clade composed of the remainder species of *Nolana* like in our previous GBSSI tree (Dillon

et al., 2007). Six clades (LFY-A, LFY-BE, LFY-C, LFY-D, LFY-F, LFY-GB, see Fig. 2) have been recovered with strong to moderate bootstrap support. Generally, the components of each clade in the *LEAFY* tree are comparable with those of species in the plastid tree except LFY-BE and LFY-GB, which are distributed from Peru to northern Chile.

4. Discussion

4.1. Monophyly of *Nolana*

Nolana was strongly supported to be closely related to the tribe Lycieae based on the plastid *ndhF* and *rbcl* sequences as well as the restriction site mapping data (Olmstead and Palmer, 1992; Olmstead and Sweere, 1994). When taxa of the tribe Lycieae were used as outgroups, the monophyly of *Nolana* was strongly supported by the plastid DNA, *LEAFY* and GBSSI data with bootstrap support of 100. Taxa of *Nolana* share the unique morphological synapomorphy of having one-seeded mericarps in the fruits.

4.2. Plastid phylogeny

Members from the cp-I clade in the plastid tree have largely overlapping distributions, with the majority confined to northern Chile (18°S–30°S) and, one species, *N. paradoxa*, extending from central to southern Chile (29°15'S–42°30'S). The cp-A clade can be easily diagnosed by a set of characters including herbs with a basal rosette of leaves, and >10 mericarps. The monophyly of the cp-A clade was also suggested in previous molecular studies of ITS and *matK* (Tago-Nakawaza and Dillon, 1999) as well as GBSSI (Dillon et al., 2007). *Nolana balsamiflua* shared a strongly supported sister relationship with *N. linearifolia*. However, this relationship is not congruent with their morphology. The non-apical style, mericarp morphology and number (~10), and weakly lignified perennial herbaceous habit of *N. linearifolia* make it easily distinguishable from *N. balsamiflua*, which is more similar to other Chilean species, e.g., *N. rostrata*, *N. filifolia* and *N. stenophylla*. The sister relationship of the *N. balsamiflua*–*N. linearifolia* clade to *N. sessiliflora* should be re-examined due to the low bootstrap support value.

The species that make up the cp-B clade are generally similar as they are herbs with showy blue to purple corollas, and ~5 mericarps in the fruits. Most species in this subclade are restricted to the Peruvian coast, 7°S–16°S. The only exceptions to the coastal distribution are *N. urubambae*, *N. lezamae* and *N. chapiensis* which occur above 2000 m and 50–500 km from the coast. However, this clade lacks internal resolution and forms a large polytomy, with the only sister relationship between *Nolana gayana*–*N. humifusa* is detected (BP = 81, PP = 100). These two species have overlapping distributions between 8°S and 15°S.

The cp-E clade contains *Nolana galapagensis* from the Galápagos Island and *N. adansonii* from southern Peru and northern Chile. Nevertheless, support for this clade is low and their relationships should be viewed with caution.

The cp-G clade has moderate support and contains species restricted to northern Chile (i.e., *N. intonsa* and *N. tarapacana*) or southern Peru (the remaining species) except *N. lycioides*, which occurs in both countries. In this clade, *Nolana inflata*, *N. weissiana* and *N. plicata* form a subclade with high bootstrap support sister to the remaining taxa that form a strongly supported subclade. In the latter subclade, the two Chilean species, *N. intonsa* and *N. tarapacana* are nested in the clades of taxa from Peru. The grouping of *N. tarapacana* with *N. arequipensis* and *N. tomentella* is weakly supported, whereas the sister relationship of *N. intonsa* with the remaining Peruvian species is strongly supported.

The cp-C clade is moderately supported and is a morphologically well-diagnosed group with woody or shrubby habit, linear leaves, large showy flowers and a Chilean distribution. These species share the synapomorphy of fused mericarps with apical stigmas. Johnston (1936) recognized this group as the segregate genus *Alona*. It has been accepted at the subgeneric level by modern workers (e.g., Tago-Nakawaza and Dillon, 1999). Two additional species, *N. balsamiflua* and *N. stenophylla* also share the fruit morphology and were included in *Alona* by Johnston (1936).

However neither species is grouped with the cp-C clade in the plastid phylogeny (Fig. 1). Rather, they are sister to the cp-A clade.

Sister to the cp-C clade is the cp-FH clade with moderate support. Within this clade, *Nolana aplocaryoides* and *N. clivicola* diverged first with the remaining taxa forming a large polytomy. Taxa of the cp-FH clade are restricted to northern Chile (22°S–30°S), either inhabiting highly saline beach dunes (e.g., *N. aplocaryoides*, *N. crassulifolia*, *N. salsoloides*, *N. peruviana* and *N. divaricata*) or occurring in inland/upland habitats (e.g., *N. leptophylla*, *N. flaccida*, *N. mollis*, *N. glauca* and *N. werdermannii*). Many species within this clade, including *N. villosa* and *N. incana*, grow in the habitats known as “aguadas” which are moist areas fed by the underground water in an otherwise dry and saline quebradas (Tago-Nakawaza and Dillon, 1999). Lack of resolution within this subclade makes it unsuitable to explain the relationships between species based on the maternally inherited plastid DNA data in this study.

4.3. *LEAFY* phylogeny

Different clones of *N. sessiliflora* form a clade (BP = 100/100) sister to the large clade consisting of the remaining *Nolana* species. The LFY-A clade contains *N. parviflora*, *N. pterocarpa*, *N. baccata*, *N. rupicola* and *N. paradoxa* with strong support. This clade is also supported by the GBSSI and plastid DNA data, and the morphology.

The LFY-D clade consists of *N. balsamiflua* and *N. stenophylla* with weak support and is sister to a strongly supported the LFY-C clade, which contains *N. rostrata*, *N. filifolia*, *N. coelestis* and *N. carnosa* (i.e., subgenus *Alona*). These two clades were within the same large clade in the previous GBSSI tree (Dillon et al., 2007), but formed a polytomy. In the plastid DNA tree, *N. balsamiflua* groups with the basal *N. sessiliflora* whereas *N. stenophylla* is nested within the cp-FH clade, which is sister to the cp-C clade. However, both clades in the plastid DNA tree are only weakly supported. The relationship between LFY-C and LFY-D are consistent with distributional patterns and morphological characters of this group, since species in both clades are restricted to northern Chile and can be diagnosed morphologically by woody or shrubby habit, large showy flowers and highly fused mericarps with apical stigmas.

The LFY-BE clade (BP = 99) includes species from southern Peru and the Galápagos Island (*N. galapagensis*). *N. galapagensis* grouped with *N. arenicola* in the GBSSI tree with weak support (Dillon et al., 2007), yet these two morphologically highly distinct species do not form a clade in the *LEAFY* tree. *N. adansonii* was closely related to *N. galapagensis* in the GBSSI tree (BP = 83) and in the plastid DNA tree (BP = 56, PP < 95), but it is sister to *N. thinophila* in the *LEAFY* tree. *N. galapagensis* and *N. adansonii* differ significantly in habit, leaves, floral structure and mericarp number. All of these taxa are southern Peruvian in distribution, only with the exception of *N. galapagensis*.

For the remaining species, the Chilean *N. clivicola* is sister to a clade (BP < 50) containing species from Chile as well as Peru. Considering the low bootstrap support, the position of *N. clivicola* needs to be further tested. The sister relationship between LFY-GB and LFY-FH also needs further study due to low support values. The LFY-FH clade is moderately supported and the taxa recovered are all distributed from northern to north-central Chile and share morphological characters including erect shrubby habit, small tubular and often white corollas and generally 5–7 mericarps. However, the bootstrap support of the internal node is generally low and the relationship among species within this clade remains unresolved.

Species in the LFY-GB clade range from central Peru to northern Chile. A subclade consisting of *N. thinophila* and different clones of *N. adansonii* is sister to the remaining species in the LFY-GB clade. The morphology and habitats of these two species are quite different. The former species forms large (>1 m in diameter) prostrate mats on near-ocean beaches and have cylindrical or terete leaves, whereas the latter species occurs at greater distances from the

ocean and has distinctly petiolate leaves. Their relationship has moderate support from the base-substitution data set, and has a high PP value and high bootstrap value when the indels are included in the analysis. Sequences from additional nuclear genes may test this relationship. Although there is strong support for the remainder clade, the internal nodes lack strong support, except for some terminal clades, such as, *N. humifusa*, *N. gayana* and *N. urubambae* and *N. cerrateana*, *N. pallida*, *N. arequipensis* and *N. tomentella* and a weakly supported clade of *N. confinis* and one clone of *N. pallida* (5c).

4.4. Reticulate evolution, lineage sorting or gene duplication

The congruence of topologies from plastid DNA and *LEAFY* data has been detected in several clades of *Nolana*. Two clades had similar or identical component taxa on both plastid DNA and *LEAFY* trees. One clade comprised of *Nolana elegans*, *N. acuminata*, *N. baccata*, *N. paradoxa*, *N. parviflora*, *N. pterocarpa* and *N. rupicola* and the other contained *N. carnosa*, *N. coelestis*, *N. rostrata* and *N. filifolia*, although the position of the two clades was not the same in the plastid DNA and the *LEAFY* trees. Morphologically species in each of these two clades are similar overall and form cohesive, well-diagnosed species groups.

Nevertheless, some relationships are more complex and not congruent among different gene trees. Strong incongruence among gene trees may be the result of processes such as reticulate evolution (especially hybridization and introgression), recombination or lineage sorting (Wendel and Doyle, 1998). A striking case is *N. sessiliflora*. It groups with the Chilean species of the clade consisting of the cp-A clade and the cp-D clade in the plastid DNA tree, whereas it has been suggested to be the first diverged species in *Nolana* in the nuclear data (GBSSI and *LEAFY*). Morphologically, it is quite distinct from taxa in the cp-D clade and very different from those of the cp-A clade. The incongruence may suggest reticulate evolution of *N. sessiliflora* with perhaps the common ancestor of the cp-D clade or the ancestor of the cp-D and the cp-A clades.

Another major incongruence concerns the subgenus *Alona* (Johnston, 1936). We sampled six of the 13 species of this subgenus. In the *LEAFY* tree, the monophyly of *Alona* was strongly supported. The *Alona* group is morphologically unique with fruits having fused mericarps and apical styles. In the plastid DNA tree (Fig. 1), species of *Alona* are in three different clades: the cp-C clade, the cp-D clade and *N. stenophylla* within the cp-FH clade. Given that species from both the cp-A and the cp-FH clades (see Fig. 1) overlap in distribution with those of *Alona*, reticulate evolution among taxa of this subgenus and the other Chilean species in the cp-A and the cp-FH clades may have occurred.

The last major incongruence between plastid DNA and *LEAFY* trees is the relationships among the Peruvian species. Both plastid DNA and *LEAFY* sequences suggest that the Peruvian species are derived and they are nested within the Chilean species (Figs. 1 and 2). *N. gayana* and *N. humifusa* share similar distributional ranges from central to northern Peru whereas *N. aticoana* is confined to southern Peru and *N. urubambae* is found nearly 500 km from the coast at the elevation of 3000 m. These taxa are all annual to perennial herbs with blue to lavender corollas. Of these taxa, only *N. gayana* has stellate pubescence and a different calyx form. In the plastid tree, *N. humifusa* and *N. gayana* group together (BP = 81) and are nested within the cp-B clade, which comprises *N. scaposae*, *N. lezamae*, *N. laxa*, *N. chapiensis*, *Nalana chancoana* and *Nalana aticoana*. In the *LEAFY* tree, they group with species mostly from the cp-G clade instead of cp-B clade in the plastid DNA tree. *Nolana inflata*, *Nolana plicata* and *N. weissiana* group together in the plastid DNA tree and are sister to the cp-G clade. They are however, sister to the LFY-BE clade in the *LEAFY* tree, which are generally corresponding to the

cp-B clade. Reticulate evolution is perhaps the most likely reason for this incongruence.

The artificial hybrids of *Nolana* (Freyre et al., 2005; Saunders, 1934) demonstrated that cross were successful between species such as far related *N. paradoxa* and *N. aplocaryoides*. These results may indirectly suggest the probability of reticulate evolution in the diversification of *Nolana*. Nevertheless, lineage sorting (especially in the plastid DNA phylogeny) can not be ruled out because the branches in the plastid DNA tree are comparatively short and some of the conflicting clades are only weakly or moderately supported. But this interpretation of lineage sorting is hampered by the general lack of informative sites in the plastid genome at the species level.

We detected major incongruence between the phylogeny of GBSSI and the other two markers (plastid and *LEAFY*) concerning the two large clades in the GBSSI tree, each containing elements from both Chile and Peru. The two major clades in the GBSSI tree each also exhibit a high level of morphological diversity, yet they are strongly supported with high bootstrap values and each has a branch length much longer than most other terminal branches (ML tree not shown). The results from the parsimony and likelihood analyses of GBSSI data are congruent, suggesting that long-branch attraction is unlikely (Sanderson et al., 2000) for most branches with perhaps the exception of the *N. adansonii*–*N. galapagensis* clade. Lineage sorting due to ancient polymorphisms of the same orthologous gene copy may also be ruled out because of the long internal branches of these two major clades. An alternative hypothesis of gene duplication of the GBSSI gene may be reasonable for explaining this incongruence. However, this hypothesis is not consistent with (1) the absence of direct evidence that two or more copies from the same sample and (2) some morphologically cohesive species (e.g., species of the cp-A or LFY-A clade and of the subgenus *Alona*) grouping together instead of randomly resolving into both major clades. Because only five samples were cloned and no more than 20 clones were sequenced in our study, inadequate sampling of clones may have not recovered all copies. The second situation may be refuted if PCR selection occurs, i.e., the reaction favored certain paralogs of a multicopy gene because of differences in primer affinity related to differences in primary or secondary structure of DNA at the potential target sites (Wagner et al., 1994). Moreover, Lynch and Conery (2000) estimated an average half-life of duplicate gene copies to be about 4-million years. GBSSI was recognized as a single-copy gene in many plant families, but duplicated GBSSI copies may be undetected in some previous studies due to insufficient sampling of species and genomes. A particularly compelling example of this situation is the studies of GBSSI for *Spartina* (Poaceae) (Baumel et al., 2002; Fortune et al., 2007). Baumel et al (2002) initially detected only one copy of the GBSSI gene for most species of the *Spartina*. A further study with more clones sampled revealed repeated gene duplication followed by deletion or sometimes without deletion (Fortune et al., 2007). In the case of *Nolana*, duplication of the GBSSI gene may have occurred in the early history of the genus, and we perhaps have two main copies of the gene in *Nolana* corresponding to the two major clades (clade I and clade II in Dillon et al. (2007)).

4.5. Implications on biogeographic diversification

The *LEAFY* data suggested the basal-most position of the Chilean *N. sessiliflora*. The Chilean *N. acuminata* group (LFY-A) and the *Alona* group (LFY-C and LFY-D) then diverged next. Even though the basal-most position of *N. sessiliflora* was not detected in the plastid DNA phylogeny, it is nested within the clade of the genus consisting of the *N. acuminata* group and the *Alona* group from Chile. Reticulate evolution may have complicated the construction of the early

diversification history of the basally branching taxa or their ancestors. Nevertheless, our *LEAFY* data suggest the basal position occupied by taxa from Chile and all Peruvian species are supported to be nested within groups of Chilean taxa.

There are at least two cases of secondary dispersal/migration from Peru to Chile on the species level. The northern Chilean species *N. intonsa* is nested within a clade of Peruvian species in both plastid DNA and the nuclear trees (*LEAFY* and GBSSI), suggesting its dispersal/migration from Peru to northern Chile. *Nolana intonsa* is also morphologically similar to *N. lycioides*, *N. cerrateana* and *N. pallida* from Peru. Another case is the northern Chilean species *N. tarapacana*, which is nested in a clade of Peruvian species in the plastid DNA tree. However, the *LEAFY* sequences of *N. tarapacana* were not available and it formed a polytomy with other species from Peru in the GBSSI tree (Dillon et al., 2007).

The GBSSI data suggested a close relationship between *N. galapagensis* from the Galápagos Islands and the Peruvian *N. adansonii* and *N. arenicola* (Dillon et al., 2007). Morphologically, *N. galapagensis* is similar to the Chilean *N. sedifolia* in a set of characters including the robust shrub habit, succulent leaves and small white tubular corollas. In the plastid DNA tree, *N. galapagensis* is sister to *N. adansonii* from Peru with weak support and the Peruvian *N. arenicola* groups other Peruvian species (BP = 100). In the *LEAFY* tree *N. galapagensis* is nested in a group of Peruvian species including *N. arenicola* along with a few other Peruvian species (clade LFY-BE in Fig. 2). Although the position of *N. galapagensis* needs to be further resolved, our results support the evolution of *N. galapagensis* of its Peruvian relatives. Its morphological similarities with the Chilean *N. sedifolia* may be due to convergence or adaptive evolution after it reached the Galápagos Islands.

4.6. SINE or SINE-like insertions in *Nolana*

In recent years, a new source of phylogenetic characters, transposable elements, especially SINE (short interspersed repetitive element) families, have been employed as a unique tool for phylogenetic study (Ray, 2007; Shedlock and Okada, 2000). The utility of SINE has been basically restricted to animal phylogenetic reconstruction (Lum et al., 2000; Murata et al., 1993; Nikaido et al., 2006, 2007; Shimamura et al., 1997) and has not attracted much attention among plant phylogenetics. Only a few SINES have been employed as phylogenetic markers in plants, including the SINE detected in GBSSI exclusively in the monophyletic tribe of Hyoscyameae (Yuan et al., 2006), a putative relative of *Nolana*. At least one of these insertions from the *LEAFY* second intron may be identified as a SINE. This insertion (labeled as SINE1 in Fig. 2) is about 789 bp in size between the position 3119 and the position 3908 in the alignment and is flanked by a repeat of AATC-CAAAAT. The SINE1 occurs exclusively in a strongly supported clade of species from Chile and can be aligned with the TS (Tobacco SINE) sequences. The TTG repeat of variable length at the 3' end of the SINE1 sequence was considered to be characteristic of the TS family (Yoshioka et al., 1993). The second SINE-like insertion was detected exclusively for a clade (BP = 99/100) of species from Peru and Chile (labeled SINE2 in Fig. 2). This SINE-like insertion is ca. 472 bp in size and is flanked by a repeat of GGWGT. The third SINE-like insertion was detected exclusively for *N. paradoxa*–*N. rupicola* clade. It is about 263 bp in size and is flanked by a sequence repeat of ACTAGRAAT. Two additional SINE-like insertions were found in *N. werdermannii* (512 bp flanked by TTTAGTT) and *N. aplocaryoides* (218 bp flanked by ASCCTS), respectively. All these five insertions are longer than 200 bases and are flanked by a short direct repeat of sequences, which have been considered a hallmark of transposition and

retroposition (Li, 1997). The three SINES possessed by the three clades (SINE1, SINE2 and SINE3) corroborate the monophyly of these clades, supporting the significance of the SINES in phylogeny reconstruction. The later two SINE-like insertions (SINE4 and SINE5) are only autapomorphies for each of the two species (*N. aplocaryoides* and *N. werdermannii*). The functions of the SINES or SINE-like insertions in *Nolana* need to be explored and may be helpful for understanding the molecular evolution of the *LEAFY* gene in the genus.

Acknowledgments

Support of the study was provided by National 973 Project of China (2007CB411601), National Science Foundation of China (30625004), the National Science Foundation (DEB-0415573 to Michael Dillon and Jun Wen) and the Laboratory of Analytical Biology at the National Museum of Natural History of the Smithsonian Institution. We thank the many colleagues who have helped with field and lab work, discussed the study and/or supplied plant material for this study, including: Federico Luebert, Jeff Hunt, Lee Weigt, Zelong Nie, Kin Han, Nicolás García, Marucha Benavides, Magda Chanco, Diane Dillon, Miguel and Erica Finger, Jorge Guerra, Oswaldo Ibicache, Mauricio Ibicache, Paulina Ossa, Manuel and Gabriela Ossa, José Miguel Fariña, Segundo Leiva, Pedro Lezama, Victor Quipuscoa, Eduardo and Tina Martell, Rachel Pinto, Adriana Hoffmann, Clodomiro Marticorena, Max Quezada, Eric Rodriguez, Abundio Sagástegui, Sebastián Teillier, Carmen Trujillo Cadiz, Alfredo Tupuachi, Maximino (Nino) Villarroel, Mario Zapata, Miyuki Nakazawa, Mélica Muñoz-Schick, Sergio Moreira, Aldo Mesa, Mikio Ono, Rosanna Freyre, Edgardo Ortiz, Massiel Corrales, Gina Castillo, Juan Gonzales and Gina Arancio.

References

- Aagaard, J.E., Olmstead, R.G., Willis, J.H., Phillips, P.C., 2005. Duplication of floral regulatory genes in the Lamiales. *Am. J. Bot.* 92, 1284–1293.
- Aagaard, J.E., Willis, J.H., Phillips, P.C., 2006. Relaxed selection among duplicate floral regulatory genes in Lamiales. *J. Mol. Evol.* 63, 493–503.
- Archambault, A., Bruneau, A., 2004. Phylogenetic utility of the *LEAFY/FLORICAULA* gene in the Caesalpinioideae (Leguminosae): gene duplication and a novel insertion. *Syst. Bot.* 29, 609–626.
- Baum, D.A., Yoon, H.S., Oldham, R.L., 2005. Molecular evolution of the transcription factor *LEAFY* in Brassicaceae. *Mol. Phylogenet. Evol.* 37, 1–14.
- Baumel, A., Ainouche, M.L., Bayer, R.J., Ainouche, A.K., Misset, M.T., 2002. Molecular phylogeny of hybridizing species from the genus *Spartina* Schreb. (Poaceae). *Mol. Phylogenet. Evol.* 22, 303–314.
- Blazquez, M.A., 1997. Illuminating flowers: Constans induces *LEAFY* expression. *BioEssays* 19, 277–279.
- Blazquez, M.A., Soowal, L.N., Lee, I., Weigel, D., 1997. *LEAFY* expression and flower initiation in *Arabidopsis*. *Development* 124, 3835–3844.
- Bombliès, K., Wang, R.L., Ambrose, B.A., Schmidt, R.J., Meeley, R.B., Doebley, J., 2003. Duplicate *FLORICAULA/LEAFY* homologs *zfl1* and *zfl2* control inflorescence architecture and flower patterning in maize. *Development* 130, 2385–2395.
- Bondeson, W.E., 1896. Gynoecial morphology and funicular germination plugs in the Nolanaceae. *Nord. J. Bot.* 6, 183–198.
- Coen, E.S., Romero, J.M., Doyle, S., Elliott, R., Murphy, G., Carpenter, R., 1990. *FLORICAULA*—a homeotic gene required for flower development in *Antirrhinum majus*. *Cell* 63, 1311–1322.
- Cronk, Q.C.B., 2001. Plant evolution and development in a post-genomic context. *Nat. Rev. Genet.* 2, 607–619.
- Cronquist, A., 1981. *An Integrated System of Classification of the Angiosperms*. Columbia University Press, New York.
- D'Arcy, W.G., 1979. The classification of the Solanaceae. In: Hawkes, J.G., Lester, R.N., Skelding, A.D. (Eds.), *The Biology and Taxonomy of the Solanaceae*. Academic Press, London, pp. 3–48.
- D'Arcy, W.G., 1991. The Solanaceae since 1976, with a review of its biogeography. In: Hawkes, J.G., Lester, R.N., Nee, M., Estrada, N. (Eds.), *Solanaceae III: Taxonomy, Chemistry, Evolution*. Royal Botanic Gardens, Kew.
- Dahlgren, R.M.T., 1980. A revised system of classification of the angiosperms. *Bot. J. Linn. Soc.* 80, 91–124.
- Dillon, M.O., Tu, T., Soejima, A., Yi, T., Nie, Z., Tye, A., Wen, J., 2007. Phylogeny of *Nolana* (Nolaneae, Solanoideae, Solanaceae) as inferred from granule-bound starch synthase I (GBSSI) sequences. *Taxon* 56, 1000–1011.

- Don, G., 1838. Solanaceae. A General History of Dichlamydeous Plants. Rivington, London. pp. 397–488.
- Evans, R.C., Alice, L.A., Campbell, C.S., Kellogg, E.A., Dickinson, T.A., 2000. The granule-bound starch synthase (GBSSI) gene in the Rosaceae: multiple loci and phylogenetic utility. *Mol. Phylogenet. Evol.* 17, 388–400.
- Felsenstein, J., 1985. Confidence-limits on phylogenies—an approach using the bootstrap. *Evolution* 39, 783–791.
- Fortune, P.M., Schierenbeck, K.A., Ainouche, A.K., Jacquemin, J., Wendel, J.F., Ainouche, M.L., 2007. Evolutionary dynamics of *waxy* and the origin of hexaploid *Spartina* species (Poaceae). *Mol. Phylogenet. Evol.* 43, 1040–1055.
- Freyer, R., Douglas, A.C., Dillon, M.O., 2005. Artificial hybridizations in five species of Chilean *Nolana* (Solanaceae). *Hortscience* 40, 532–536.
- Frohlich, M.W., Meyerowitz, E.M., 1997. The search for flower homeotic gene homologs in basal angiosperms and Gnetales: a potential new source of data on the evolutionary origin of flowers. *Int. J. Plant Sci.* 158, S131–S142.
- Frohlich, M.W., Parker, D.S., 2000. The mostly male theory of flower evolutionary origins: from genes to fossils. *Syst. Bot.* 25, 155–170.
- Gene Codes Corporation, 2005. Sequencher 4.5. Gene Codes Corporation, Ann Arbor, Michigan.
- Grob, G.B.J., Gravendeel, B., Eurlings, M.C.M., 2004. Potential phylogenetic utility of the nuclear *FLORICAULA/LEAFY* second intron: comparison with three chloroplast DNA regions in *Amorphophallus* (Araceae). *Mol. Phylogenet. Evol.* 30, 13–23.
- Hofer, J., Turner, L., Hellens, R., Ambrose, M., Matthews, P., Michael, A., Ellis, N., 1997. UNIFOLIATA regulates leaf and flower morphogenesis in pea. *Curr. Biol.* 7, 581–587.
- Hoot, S.B., Taylor, W.C., 2001. The utility of nuclear ITS, a *LEAFY* homolog intron, and chloroplast *atpB-rbcL* spacer region data in phylogenetic analyses and species delimitation in *Isoetes*. *Am. Fern J.* 91, 166–177.
- Howarth, D.G., Baum, D.A., 2005. Genealogical evidence of homoploid hybrid speciation in an adaptive radiation of *Scaevola* (Goodeniaceae) in the Hawaiian Islands. *Evolution* 59, 948–961.
- Hunziker, A.T., 2001. Genera Solanacearum. The Genera of Solanaceae Illustrated According to a New System. A.R.G. Gantner Verlag K.-G. Ruggell, Liechtenstein.
- Johnston, J.S., 1936. A study of the Nolanaceae. *Contr. Gray Herb.* 112, 1–87.
- Lynch, M., Conery, J.S., 2000. The evolutionary fate and consequences of duplicate genes. *Science* 290, 1151–1155.
- Kelly, A.J., Bonnländer, M.B., Meekswagner, D.R., 1995. NFI, the tobacco homolog of *FLORICAULA* and *LEAFY*, is transcriptionally expressed in both vegetative and floral meristems. *Plant Cell* 7, 225–234.
- Knapp, S., 2002. Tobacco to tomatoes: a phylogenetic perspective on fruit diversity in the Solanaceae. *J. Exp. Bot.* 53, 2001–2022.
- Lee, C., Wen, J., 2004. Phylogeny of *Panax* using chloroplast *trnC-trnD* intergenic region and the utility of *trnC-trnD* in interspecific studies of plants. *Mol. Phylogenet. Evol.* 31, 894–903.
- Levin, R.A., Miller, J.S., 2005. Relationships within tribe Lycieae (Solanaceae): paraphyly of *Lycium* and multiple origins of gender dimorphism. *Am. J. Bot.* 92, 2044–2053.
- Levin, R.A., Myers, N.R., Bohs, L., 2006. Phylogenetic relationships among the “Spiny Solanums” (*Solanum* subgenus *Leptostemonum*, Solanaceae). *Am. J. Bot.* 93, 157–169.
- Levin, R.A., Watson, K., Bohs, L., 2005. A four-gene study of evolutionary relationships in *Solanum* section *Acantophora*. *Am. J. Bot.* 92, 603–612.
- Li, W.-H., 1997. Molecular Evolution. Sinauer Associates, Inc., Sunderland, MA.
- Lorenz-Lemke, A.P., Mader, G., Muschner, V.C., Stehmann, J.R., Bonatto, S.L., Salzano, F.M., Freitas, L.B., 2006. Diversity and natural hybridization in a highly endemic species of *Petunia* (Solanaceae): a molecular and ecological analysis. *Mol. Ecol.* 15, 4487–4497.
- Lum, J.K., Nikaido, M., Shimamura, M., Shimodaira, H., Shedlock, A.M., Okada, N., Hasegawa, M., 2000. Consistency of SINE insertion topology and flanking sequence tree: quantifying relationships among Cetartiodactyls. *Mol. Biol. Evol.* 17, 1417–1424.
- Mesa, A., 1981. Nolanaceae. *Flora Neotrop.* 26, 1–197.
- Murata, S., Takasaki, N., Saitoh, M., Okada, N., 1993. Determination of the phylogenetic-relationships among Pacific Salmonids by using short interspersed elements (SINEs) as temporal landmarks of evolution. *Proc. Natl. Acad. Sci. USA* 90, 6995–6999.
- Nikaido, M., Hamilton, H., Makino, H., Sasaki, T., Takahashi, K., Goto, M., Kanda, N., Pastene, L.A., Okada, N., 2006. Baleen whale phylogeny and a past extensive radiation event revealed by SINE insertion analysis. *Mol. Biol. Evol.* 23, 866–873.
- Nikaido, M., Piskurek, O., Okada, N., 2007. Toothed whale monophyly reassessed by SINE insertion analysis: the absence of lineage sorting effects suggests a small population of a common ancestral species. *Mol. Phylogenet. Evol.* 43, 216–224.
- Oh, S.H., Potter, D., 2003. Phylogenetic utility of the second intron of *LEAFY* in *Neillia* and *Stephanandra* (Rosaceae) and implications for the origin of *Stephanandra*. *Mol. Phylogenet. Evol.* 29, 203–215.
- Oh, S.H., Potter, D., 2005. Molecular phylogenetic systematics and biogeography of tribe *Neillieae* (Rosaceae) using DNA sequences of cpDNA, rDNA, and *LEAFY*. *Am. J. Bot.* 92, 179–192.
- Olmstead, R.G., Palmer, J.D., 1992. A chloroplast DNA phylogeny of the Solanaceae subfamilial relationships and character evolution. *Ann. MO Bot. Gard.* 79, 346–360.
- Olmstead, R.G., Sweere, J.A., 1994. Combining data in phylogenetic systematics: an empirical approach using three molecular data sets in the Solanaceae. *Syst. Biol.* 43, 467–481.
- Olmstead, R.G., Sweere, J.A., Spangler, R.E., Bohs, L., Palmer, J.D., 1999. Phylogeny and provisional classification of the Solanaceae based on chloroplast DNA. In: Nee, M., Symon, D.E., Lester, R.N., Jessop, J.P. (Eds.), *Solanaceae IV: Advances in Biology and Utilization*. Royal Botanic Gardens, Kew, Richmond, pp. 111–137.
- Peralta, I.E., Spooner, D.M., 2001. Granule-bound starch synthase I (GBSSI) gene phylogeny of wild tomatoes (*Solanum* L. Section *Lycopersicon* (mill.) Wettst. Subsection *Lycopersicon*). *Am. J. Bot.* 88, 1888–1902.
- Perez, F., Arroyo, M.T.K., Medel, R., Hershkovitz, M.A., 2006. Ancestral reconstruction of flower morphology and pollination systems in *Schizanthus* (Solanaceae). *Am. J. Bot.* 93, 1029–1038.
- Posada, D., Buckley, T.R., 2004. Model selection and model averaging in phylogenetics: advantages of akaike information criterion and Bayesian approaches over likelihood ratio tests. *Syst. Biol.* 53, 793–808.
- Posada, D., Crandall, K.A., 1998. Modeltest: testing the model of DNA substitution. *Bioinformatics* 14, 817–818.
- Pouteau, S., Nicholls, D., Tooke, F., Coen, E., Battey, N., 1997. The induction and maintenance of flowering in *impatiens*. *Development* 124, 3343–3351.
- Rambaut, A., 2007. Se-al version 2.0a11. Available from: <<http://tree.bio.ed.ac.uk/software/seal/>>.
- Rannala, B., Yang, Z.H., 1996. Probability distribution of molecular evolutionary trees: a new method of phylogenetic inference. *J. Mol. Evol.* 43, 304–311.
- Ray, D.A., 2007. Sines of progress: mobile element applications to molecular ecology. *Mol. Ecol.* 16, 19–33.
- Ronquist, F., Huelsenbeck, J.P., 2003. MrBayes 3: Bayesian phylogenetic inference under mixed models. *Bioinformatics* 19, 1572–1574.
- Sanderson, M.J., Wojciechowski, M.F., Hu, J.M., Khan, T.S., Brady, S.G., 2000. Error, bias, and long-branch attraction in data for two chloroplast photosystem genes in seed plants. *Mol. Biol. Evol.* 17, 782–797.
- Sang, T., 2002. Utility of low-copy nuclear gene sequences in plant phylogenetics. *Crit. Rev. Biochem. Mol. Biol.* 37, 121–147.
- Saunders, E.R., 1934. The history, origin and characters of certain interspecific hybrids in *Nolana* and their relation to *Nolana paradoxa*. *J. Genet.* 29, 387–419.
- Saunders, E.R., 1936. On certain unique features of the gynoeceum in Nolanaceae. *New Phytol.* 35, 423–431.
- Schultz, E.A., Haughn, G.W., 1991. *LEAFY*, a homeotic gene that regulates inflorescence development in *Arabidopsis*. *Plant Cell* 3, 771–781.
- Shaw, J., Lickey, E.B., Beck, J.T., Farmer, S.B., Liu, W., Miller, J., Siripun, K.C., Winder, C.T., Schilling, E.E., Small, R.L., 2005. The tortoise and the hare II: relative utility of 21 noncoding chloroplast DNA sequences for phylogenetic analysis. *Am. J. Bot.* 92, 142–166.
- Shaw, J., Lickey, E.B., Schilling, E.E., Small, R.L., 2007. Comparison of whole chloroplast genome sequences to choose noncoding regions for phylogenetic studies in angiosperms: the tortoise and the hare III. *Am. J. Bot.* 94, 275–288.
- Shedlock, A.M., Okada, N., 2000. SINE insertions: powerful tools for molecular systematics. *BioEssays* 22, 148–160.
- Shimamura, M., Yasue, H., Ohshima, K., Abe, H., Kato, H., Kishiro, T., Goto, M., Munechika, I., Okada, N., 1997. Molecular evidence from retrotransposons that whales form a clade within even-toed ungulates. *Nature* 388, 666–670.
- Smith, S.D., Baum, D.A., 2006. Phylogenetics of the florally diverse andean clade Iochrominae (Solanaceae). *Am. J. Bot.* 93, 1140–1153.
- Souer, E., van der Krol, A., Kloos, D., Spelt, C., Blied, M., Mol, J., Koes, R., 1998. Genetic control of branching pattern and floral identity during *Petunia* inflorescence development. *Development* 125, 733–742.
- Southerton, S.G., Strauss, S.H., Olive, M.R., Harcourt, R.L., Decroocq, V., Zhu, X.M., Llewellyn, D.J., Peacock, W.J., Dennis, E.S., 1998. *Eucalyptus* has a functional equivalent of the *Arabidopsis* floral meristem identity gene *LEAFY*. *Plant Mol. Biol.* 37, 897–910.
- Tago-Nakawaza, M., Dillon, M.O., 1999. Biogeografía y evolución en el clado *Nolana* (Solanaceae-Solanaceae). *Arnaldoa* 6, 81–116.
- Takhtajan, A.L., 1997. Diversity and Classification of Flowering Plants. Columbia University Press, New York.
- Theissen, G., 2000. Plant breedings: FLO-like meristem identity genes: from basic science to crop plant design. *Progress in botany*. Springer-Verlag, Berlin. pp. 167–183.
- Thompson, J.D., Gibson, T.J., Plewniak, F., Jeanmougin, F., Higgins, D.G., 1997. The Clustal windows interface. Flexible strategies for multiple sequence alignment aided by quality analysis tools. *Nucleic Acids Res.* 25, 4876–4882.
- Thorne, R.F., 1983. Proposed new realignments in the angiosperms. *Nord. J. Bot.* 3, 85–117.
- Wada, M., Cao, Q.F., Kotoda, N., Soejima, J., Masuda, T., 2002. Apple has two orthologues of *FLORICAULA/LEAFY* involved in flowering. *Plant Mol. Biol.* 49, 567–577.
- Wagner, A., Blackstone, N., Cartwright, P., Dick, M., Misof, B., Snow, P., Wagner, G.P., Bartels, J., Murtha, M., Pendleton, J., 1994. Surveys of gene families using polymerase chain reaction: PCR selection and PCR drift. *Syst. Biol.* 43, 250–261.
- Weigel, D., 1995. The genetics of flower development: from floral induction to ovule morphogenesis. *Annu. Rev. Genet.* 29, 19–39.
- Wendel, J.F., Doyle, J.J., 1998. Phylogenetic incongruence. Window into genome history and molecular evolution. In: Soltis, D.E., Soltis, P.S., Doyle, J.J. (Eds.), *Molecular Systematics of Plants II: DNA Sequencing*. Kluwer Academic Publishers, Boston, pp. 1–42.
- Winkworth, R.C., Donoghue, M.J., 2004. *Viburnum* phylogeny: evidence from the duplicated nuclear gene GBSSI. *Mol. Phylogenet. Evol.* 33, 109–126.
- Yoshioka, Y., Matsumoto, S., Kojima, S., Ohshima, K., Okada, N., Machida, Y., 1993. Molecular characterization of a short interspersed repetitive element from

- tobacco that exhibits sequence homology to specific transfer-RNAs. *Proc. Natl. Acad. Sci. USA* 90, 6562–6566.
- Young, N., Healy, J., 2003. Gapcoder automates the use of indel characters in phylogenetic analysis. *BMC Bioinformatics* 4, 6.
- Yuan, Y.W., Zhang, Z.Y., Chen, Z.D., Olmstead, R.G., 2006. Tracking ancient polyploids: a retroposon insertion reveals an extinct diploid ancestor in the polyploid origin of belladonna. *Mol. Biol. Evol.* 23, 2263–2267.